

Condensation and evolution of space-time network

Bi Qiao^{1,2}

¹*Department of Physics, Science School,
Wuhan University of Technology, Wuhan 430070,
China;* ²*International Noble Academy,
1075 Ellesmere Road, Toronto, M1P 5C3 Canada*

Abstract

In this work, we try to propose, in a novel way using the Bose and Fermi quantum network approach, a framework studying condensation and evolution of space time network described by the Loop quantum gravity. Considering quantum network connectivity features in the Loop quantum gravity, we introduce a link operator, and through extending the dynamical equation for the evolution of quantum network posed by Ginestra Bianconi to an operator equation, we get the solution of the link operator. This solution is relevant to the Hamiltonian of the network, and then is related to the energy distribution of network nodes. Showing that tremendous energy distribution induce huge curved space-time network, may have space time condensation in high-energy nodes. For example, in the black hole circumstances, quantum energy distribution is related to the area, thus the eigenvalues of the link operator of the nodes can be related to quantum number of area, and the eigenvectors are just the spin network states. This reveals that the degree distribution of nodes for space-time network is quantized, which can form the space-time network condensation. The black hole is a sort of result of space-time network condensation, however there may be more extensive space-time network condensation, for example, the universe singularity (big bang).

Keywords: Loop quantum gravity, Spin network, Complex network, Quantum network, Black hole.

I. INTRODUCTION

The greatest challenging topic to contemporary fundamental physics is to coordinate general relativity and quantum mechanics, quantum gravity research. This study will cause us to have profound changes in the concept on space, time and matter structure, and achieve a quantum revolution. At present, for the quite progressed theory of quantum gravity, there are two forms. First, quantized general relativity, canonical quantum gravity belongs to this. Second, quantized a classic theory which is deferent from general relativity and the general relativity is as its low-energy limit. The super-string / M theory belongs to this. Loop quantum gravity ^[1] is the current representative form of canonical quantum gravity. Canonical quantum gravity is one only when the gravitational effects of quantum gravity. Comparing with superstring / M theory it does not include other different interactions. Its basic concept is to apply standard quantization procedure to general relativity while general relativity is written in the canonical (Hamiltonian) form. According to historical development the canonical quantum gravity can be divided into two stages, the simple quantum gravity and Loop quantum gravity. Roughly speaking, the former occurred before 1986, and later occurred after 1986. In simple quantum gravity, there is renormalization difficulty for UV divergence, and Loop quantum gravity has become currently the representative.

According to basic spirit of the Loop quantum gravity space-time is the gravitational field and space-time is quantized in the Planck scale, as well formed space-time network which uses quantum of the volume as a node, connected through the quantum of adjacent area. The basis of the space time network can be described by spin network. Physical space time can be expanded by spin networks as basis. The evolution of spin network can be studied by calculating spinfoam. Spinfoam is main method to study evolution of spin network, based on the Feynman propagation in the Loop quantum gravity. The current studied complex network theoretical method ^[2] is irrelevant to the study of the space time network.

In this work, we try to use the Bose and Fermi quantum network approach to present a framework for studying condensation and evolution of space time network. Considering quantum network connectivity features, we introduce a link operator, and through extending the complex network equation for the evolution of quantum network in the literature [3] to an operator equation, we get the solution of the link operator.

This solution is relevant to the Hamiltonian of the network, and then is related to the

energy distribution of network nodes. Showing that tremendous energy distribution induce huge curved space-time network, may have space time condensation in high-energy nodes. For example, in the black hole circumstances, quantum energy distribution is related to the area, thus the eigenvalues of the degree operator of the nodes can be related to quantum number of area, and the eigenvectors are just the spin network states. This reveals that the degree distribution of nodes for space-time network is quantized, which can form the space-time network condensation. The black hole is a sort of space-time network condensation, however there may be more extensive network of space-time setting, for example, the universe singularity (big bang).

II. CONNECTIVITY OF SPIN NETWORK

Let a graph Γ to be given immersed in the manifold Σ , for which an ordering and an orientation have been chosen. Denote the nodes as the end points of the oriented curves in Γ , and joined by links l . Then j_l is an assignment of an irreducible representation to each link l and i_n is an assignment of an intertwiner to each node n . The intertwiner i_n associated with a node is between the representations associated with the links adjacent to the node. Γ becomes a collection of nodes n with links l . The triplet $S \equiv (\Gamma, i_n, j_n)$ is called a spin network embedded in Σ . The physical space-time is combination of spin networks. More precisely, j_n represents unitary irreducible representations and i_n represents a basis in the space of the intwiners between the representations adjacent to the nodes n . The area of a surface cutting n links of spin network with $l = 1, \dots, n$ quantized labels j_l , given by the eigenvalue of the area operator, $8\pi\gamma\frac{\hbar}{2\pi}G\sum_n\sqrt{j_n(j_n+1)}$, where γ is the Immirzi parameter which is a free dimensionless constant and G is the Newton constant. Consider the quantum property of the connectivity for nodes of spin network, we introduce a link (degree) operator of node $\mathbf{K}$ which acts to the spin network state $|S\rangle$ and gives the eigenvalue j_n by

$$\mathbf{K}|S\rangle = \tau^{f(j_n)}|S\rangle, \quad (1)$$

where f is a function of j_n , and $|S\rangle = (\Gamma, i_n, \alpha_n, \beta_n)$ is given by the spin network $S \equiv (\Gamma, i_n, j_n)$ in such way : $\langle U|j, \alpha, \beta\rangle = (R^j(U))_\alpha^\beta$ forms an orthonormal basis in the Hilbert space $L_2[G, dU]$, and introduces a spin network state $|S\rangle$ by the Peter-Weyl theorem^[1].

Extending the rate equation of connectivity in ref.[2] to the link operator $\mathbf{K}$, a dynamical

equation for $\mathbf{K}$ can be constructed by

$$\frac{\partial \mathbf{K}}{\partial \tau} = m_+ \frac{e^{\beta H_0} \mathbf{K}}{\text{Tr}(e^{\beta H_0} \mathbf{K})} - m_- \frac{e^{-\beta H_0} \mathbf{K}}{\text{Tr}(e^{-\beta H_0} \mathbf{K})}, \quad (2)$$

where τ is introduced as a proper time in this relativity system, the spin network state is associated with a general three-dimensional spacelike hypersurface, m_+ and m_- are proportional (suitable) coefficients (constant or function) for making the equation (2) to be held, and H_0 is an energy part of the (constrained) Hamiltonian of the system H given by

$$H = \int N \text{tr}(F \wedge \{\mathbf{V}, A\}) \quad (3)$$

$$= p_t + H_0, \quad (4)$$

here F (curvature) and A (complex field) are limits of holonomy operators of small paths, $\{\}$ is the Poisson bracket as quantum commutator, and $\mathbf{V}$ is the quantum volume operator^{[1],[5]}.

In the thermodynamical limit, $\text{Tr}(e^{\beta H_0} \mathbf{K})$ and $\text{Tr}(e^{-\beta H_0} \mathbf{K})$ are supposed to tend to

$$\text{Tr}(e^{\beta H_0} \mathbf{K}) \rightarrow m_+ \tau e^{\beta \mu_+}, \quad (5)$$

$$\text{Tr}(e^{-\beta H_0} \mathbf{K}) \rightarrow m_- \tau e^{-\beta \mu_-}, \quad (6)$$

where μ_+ , μ_- are suitable constants which are related to the asymptotic behavior of the network. This is so because $\text{Tr}(e^{\beta H_0} \mathbf{K})$ and $\text{Tr}(e^{-\beta H_0} \mathbf{K})$ can be assumed to be self-average and converge to their mean value and grow linearly in time under suitable conditions^[4].

Thus, Eq.(2) becomes

$$\frac{\partial \mathbf{K}}{\partial \tau} = \left[e^{\beta(H_0 - \mu_+)} - e^{-\beta(H_0 - \mu_-)} \right] \frac{\mathbf{K}}{\tau}, \quad (7)$$

and the solution of the equation therefore is given by

$$\mathbf{K} = \tau^{f(H_0)}, \quad (8)$$

where $f(H_0)$ is defined by

$$f(H_0) = e^{\beta(H_0 - \mu_+)} - e^{-\beta(H_0 - \mu_-)}. \quad (9)$$

This allows one to obtain the mean value $\langle \mathbf{K} \rangle$ as

$$\langle \mathbf{K} \rangle = (m_+ + m_-) \left(\frac{\tau}{\tau'} \right)^{f(M)}, \quad (10)$$

with the initial condition

$$\langle \mathbf{K} \rangle_0 = m_+ + m_-, \quad (11)$$

where M represents the eigenvalue of H_0 which is related to the energy in the nodes, and relevant tracing calculation is performed with respect to the eigen-states of H_0 . When H_0 acts on a spin network state, this operator acts only on the nodes of the spin network, because of the presence of the volume in the Hamiltonian operator H , hence $\langle \mathbf{K} \rangle$ can represent a general connectivity of space-time network, this space-time network can be expanded by many spin networks as unit networks.

III. CONDENSATION OF SPACE-TIME NETWORK

In the Schwarzschild black hole environment, the energy M is related to the area A through^[6]

$$M = \sqrt{\frac{A}{16\pi G^2}}. \quad (12)$$

Thus, the mean value $\langle \mathbf{K} \rangle$ becomes

$$\langle S | \mathbf{K} | S \rangle = (m_+ + m_-) \left(\frac{\tau}{\tau'} \right)^{f\left(\sqrt{\frac{A}{16\pi G^2}}\right)} = (m_+ + m_-) \left(\frac{\tau}{\tau'} \right)^{f\left(\sqrt{\frac{\frac{h}{2\pi}\gamma\sqrt{j_n(j_n+1)}}{2G}}\right)}, \quad (13)$$

where $f\left(\sqrt{\frac{1}{2G}\frac{h}{2\pi}\gamma\sqrt{j_n(j_n+1)}}\right)$ is a function of $\sqrt{j_n(j_n+1)}$ or j_n . This shows that $\langle S | \mathbf{K} | S \rangle$ is quantized, with quantum number of link as $\sqrt{j_n(j_n+1)}$, where j_n is determined by eigenvalues of the area operator $\mathbf{A}$ with respect to spin network state S and represents the quantum number of the area. One of properties of the quantized links of nodes is that there are possibly many links labelled by quantum numbers between two adjacent nodes with relevant quantum weights, while the classical link between the same two nodes is just one link with (or without) a classical weight.

Consequently, one can get

$$\frac{m}{m+m'} = \int dMp(M) \frac{e^{\beta(M-\mu_+)}}{1 - e^{\beta(M-\mu_+)} + e^{-\beta(M-\mu_-)}}, \quad (14)$$

and

$$\frac{m'}{m+m'} = \int dMp(M) \frac{e^{-\beta(M-\mu_+)}}{1 - e^{\beta(M-\mu_+)} + e^{-\beta(M-\mu_-)}}. \quad (15)$$

Here the ratio $\frac{m}{m+m'}$ or $\frac{m'}{m+m'}$ represents a fraction of m or m' links attached or detached to nodes in the evolution of the network. The former represents the increase of the links on the node, and the second one represents the decrease of links on the node. When $\mu_+ = \mu_- = 0$ and $M \rightarrow 0$, it is

$$\frac{m}{m+m'} = \frac{m'}{m+m'} = \int dM p(M), \quad (16)$$

which shows that the quantized connectivity of the space-time network, in the very low energy condition, evolves either increase or decrease in time as a mixed process of attached or detached to nodes. While $\mu_+ = \mu_- = 0$ and $M \rightarrow \infty$, it is not difficult to find $\frac{m'}{m+m'} = 0$, and one gets

$$\frac{m}{m+m'} = - \int dM p(M). \quad (17)$$

This shows, in this situation, there is only link increased process on the node while the detached process is vanished! This reveals that a kind of condensation of space-time network around the huge energy nodes may happen because there is only a monotonic attached link process in the evolution of the space-time network. For example, in the (Schwarzschild) black hole surroundings, there is a sort of space-time network condensation on the singularity with $M \rightarrow \infty$. Furthermore, in the universal space-time scale, there may be more extensive space-time network condensation, such as the universe condensates itself as one (many) singularity, a (many) big bang.

IV. CONNECTIVITY DISTRIBUTION

The connectivity distribution $P(\langle \mathbf{k} \rangle)$ is given by the sum of the probabilities of the nodes with energy M and connectivity $\langle \mathbf{k} \rangle$. Following the idea in ref.[2], considering the degree of the nodes in our model k changes to the mean value $\langle \mathbf{K} \rangle$ since the quantum property of the link for the space-time network, we have to sum over all the modes with energy lower or higher than a threshold defined as $\epsilon_s = \frac{\mu_+ + \mu_-}{2}$, when $\langle \mathbf{k} \rangle < m + m'$ or $\langle \mathbf{k} \rangle > m + m'$,

respectively,

$$\begin{aligned}
P(\langle \mathbf{k} \rangle) &= \theta(\langle \mathbf{k} \rangle - \langle \mathbf{k} \rangle_0) \int_{\epsilon_s > \frac{\mu_+ + \mu_-}{2}} dMp(M) \frac{1}{\frac{\partial \langle \mathbf{k} \rangle}{\partial \tau}} + \theta(\langle \mathbf{k} \rangle_0 - \langle \mathbf{k} \rangle) \int_{\epsilon_s < \frac{\mu_+ + \mu_-}{2}} dMp(M) \frac{1}{\frac{\partial \langle \mathbf{k} \rangle}{\partial \tau}} \\
&= \theta(\langle \mathbf{k} \rangle - m - m') \frac{1}{m + m'} \int_{\epsilon_s > \frac{\mu_+ + \mu_-}{2}} dMp(M) \frac{1}{f(M)} \left(\frac{\langle \mathbf{k} \rangle}{m + m'} \right)^{-1 - \frac{1}{f(M)}} \\
&\quad + \theta(m + m' - \langle \mathbf{k} \rangle) \frac{1}{m + m'} \int_{\epsilon_s < \frac{\mu_+ + \mu_-}{2}} dMp(M) \frac{1}{f(M)} \left(\frac{\langle \mathbf{k} \rangle}{m + m'} \right)^{-1 - \frac{1}{f(M)}}.
\end{aligned} \tag{18}$$

If $\mu_+ = \mu_- = 0$ and $M \rightarrow \infty$, it is possible that $\frac{1}{f(M)} = 0$, and $\int_{\epsilon_s > \frac{\mu_+ + \mu_-}{2}} dMp(M) \frac{1}{f(M)}$ is finite (convergent and $\neq 0$), which results in

$$P(\langle \mathbf{k} \rangle) = \theta(\langle \mathbf{k} \rangle - m - m') \frac{1}{m + m'} \int_{\epsilon_s > \frac{\mu_+ + \mu_-}{2}} dMp(M) \frac{1}{f(M)} \left(\frac{\langle \mathbf{k} \rangle}{m + m'} \right)^{-1} \sim B \frac{1}{\langle \mathbf{k} \rangle}, \tag{19}$$

where the coefficient B is defined by

$$B = \int_{\epsilon_s > \frac{\mu_+ + \mu_-}{2}} dMp(M) \frac{1}{f(M)}, \tag{20}$$

and the integral variable is supposed to be irrelevant to $\langle \mathbf{k} \rangle$. This result means that the connectivity distribution $P(\langle \mathbf{k} \rangle)$, in the above approximations, obeys the power law and the space-time network is scale-free, the connectivity distribution $P(\langle \mathbf{k} \rangle)$ is inversely proportional to the increase of the connectivity $\langle \mathbf{k} \rangle$. This means the probability of $\langle \mathbf{k} \rangle$ tending to infinity is quite small if value of B is tiny. However, if B is big or is a function of $\langle \mathbf{k} \rangle$, then situation of $P(\langle \mathbf{k} \rangle)$ is complicated, then, the relevant space-time network may be not scale-free.

V. CONCLUSIONS

The quantum connectivity of space-time network can be calculated by introducing the link operator. The mean value of the link operator is a function of eigenvalue for the energy part of the Hamiltonian operator of the space-time network. In the Schwarzschild black hole situation, the mean value of the link operator is a function of quantum number of the area operator with respect to the spin network states, and the connectivity is quantized. Moreover, there is a sort of condensation of space-time network around its extreme high

energy nodes. In the condensation progression there exists only a monotonic attached link process in the evolution of the network, which may results in the singularity in the universe. The connectivity distribution can be represented by Eq.(19), which may or may not obey the power law.

Acknowledgement: This work was supported by the grants from NSERC, CIPI, MMO, CITO, Wuhan University of Technology, and Chinese NSFC under the Grand No. 60874087.

-
- [1] Carlo Rovelli, *Quantum Gravity*, Cambridge University Press, (2004).
 - [2] Reka Albert, and Albert-Laszlo Barabasi, *Statistical mechanics of complex networks*, Rev. Modern Phys. 74, 47 (2002).
 - [3] Ginestra Bianconi, *Quantum statistics in complex networks*, Phys. Rev. E 66, 056123 (2002).
 - [4] Ginestra Bianconi, and Albert-Laszlo Barabasi, *Bose-Einstein condensation in complex networks*, Phys. Rev. Lett. 86, 5632 (2001).
 - [5] Thomas Thiemann, *Modern canonical quantum general relativity*, Cambridge University Press, Cambridge, New York, Melbourne, (2007).
 - [6] Charles W. Misner, Kip S. Thorne, and John Archibald Wheeler, *Gravitation*, W. H. Freeman And Company, San Francisco, (1973).